

Effective Mass of a Vortex in a Clean Superconductor

J. H. Han¹, P. Ao² and X.-M. Zhu²

Asia Pacific Center for Theoretical Physics, 207-43 Cheongryangri-dong Dongdaemun-gu Seoul 130-012, Korea¹
and

Department of Physics, Umeå University, S-901 87 Umeå, Sweden²

We calculate the effective mass of a single quantized vortex in the BCS superconductor at finite temperature. Using the effective action for a vortex, we arrive at the mass formula as the integral of the spectral function $J(\omega)/\omega^3$ over frequency. The spectral function is given in terms of the transition elements of the gradient of the Hamiltonian between Bogoliubov-deGennes eigenstates. We show that core-core, and core-extended transitions yield the vortex mass, near $T=0$, of order of electron mass displaced by the normal core. The extended-extended states contributions are linearly divergent with the low frequency cutoff ω_c , in accordance with the Ohmic character of the spectral function at low energies. We argue that the mass and friction are closely related in this system and arise from the same mechanism - interaction with the surrounding fermionic degrees of freedom.

The dynamics of an individual vortex or their array is an integral part of our understanding of the superfluid matter. In superconductors there have been phenomenological approaches to write down a classical equation of motion of a vortex [1]. This “effective” equation will in general rely on the information of the mass, along with the longitudinal and transverse forces that act on the vortex.

The vortex mass also enters in an important way in the tunneling process [2,3]. Estimates of the mass based on Ginzburg-Landau functional gives a mass of order mk_f per unit vortex length [4], while more recent work employing microscopic BCS-type analysis predicts a much larger mass $\sim mk_f(k_f\xi_0)^2$ [5,6]. The authors of Refs. [5,6] focused on the core quasiparticle contribution to mass, and were limited to zero temperature. We use in this work the effective action approach [6,7] to go beyond those limitations.

A quantized vortex is a phase singularity accompanied by a change in the order parameter density near the center. A closer scrutiny using the Bogoliubov-de Gennes (BdG) equation reveals that there are discrete energy levels inside the vortex core [8], in addition to a continuous spectrum with energy $|E|$ larger than the gap whose wavefunction extends beyond the core region. Both states are perturbed once the vortex is set in motion, by a perturbing Hamiltonian equal to the inner product of the fermion current with the vortex velocity. Virtual transitions are induced among core and extended states and lead to an energy shift proportional to the square of the vortex velocity. Dividing by the squared velocity gives a mass in agreement with those of Refs. [5,6].

Here we proceed via an effective action in the Matsubara representation. By integrating out the fermions from the BCS action and expanding the remaining degrees of freedom up to second order in the vortex coordinate one can obtain the effective action of a single vortex [6,7]. It is possible to *derive* the transverse as well as the longitudinal forces on a vortex from this action [7]. Likewise we

derive below a formal expression of the vortex mass valid at finite temperature. It agrees with the Hamiltonian perturbation theory at $T = 0$. Since core and extended states have non-vanishing overlap in the core region one expects core-core, core-extended and extended-extended transitions to contribute to the mass unlike previous theories which only considered core-core processes. Using approximate forms for eigenstates we explicitly compute the transition elements and the mass. At low temperature the mass is comparable to that of electrons displaced by the hollow core of the vortex.

The effective action by itself is non-local in time, hence dissipative, and the concept of mass only arises as a result of the local-time approximation. We have deduced both the mass and friction of the vortex from a single framework, with the relative significance of inertia and dissipation largely determined by the motion itself. In this work we will focus on the pure system with no disorder.

The effective action of a single vortex, characterized by the two-dimensional coordinate $r_v(\tau)$ [9] in the BCS superconductor, is given by [6,7]

$$S_{eff} = \frac{1}{2\pi} \int_0^\beta d\tau \int_{-\infty}^\infty d\tau' |r_v(\tau) - r_v(\tau')|^2 \times \int_0^\infty d\omega e^{-\omega|\tau-\tau'|} J(\omega). \quad (1)$$

In the theory of dissipative quantum systems $J(\omega)$ is known as the spectral function [3]:

$$J(\omega) = \frac{\pi}{4} \sum_{\alpha\beta} \delta(\omega - |E_\alpha - E_\beta|) |f(E_\alpha) - f(E_\beta)| |\langle \alpha | \nabla_v H_0 | \beta \rangle|^2. \quad (2)$$

It is the gradient of the Hamiltonian operator that governs the longitudinal dynamics of the vortex. The transverse motion of the vortex, which is of course crucial for the overall dynamics, is not explicitly considered in this work. The longitudinal action shown in Eq. (1) is responsible for the mass and also dissipation of the vortex. The

states α, β are the eigenstates of the Nambu Hamiltonian H_0 with the energy E_α and E_β respectively.

By assuming $r_v(\tau) - r_v(\tau') \approx (\tau - \tau')\dot{r}_v(\tau)$, one can rewrite the action as

$$S_{eff} \approx \frac{1}{2} \int_0^\beta d\tau \left\{ \int_0^\infty d\omega \frac{4J(\omega)}{\pi\omega^3} \right\} \left(\frac{dr_v}{d\tau} \right)^2. \quad (3)$$

The vortex mass follows as

$$M_v = \sum_{\alpha\beta} \left| \frac{f(E_\alpha) - f(E_\beta)}{(E_\alpha - E_\beta)^3} \right| |\langle \alpha | \nabla_v H_0 | \beta \rangle|^2 \quad (4)$$

$$= \sum_{\alpha\beta} \left| \frac{f(E_\alpha) - f(E_\beta)}{E_\alpha - E_\beta} \right| |\langle \alpha | \nabla_v | \beta \rangle|^2. \quad (5)$$

for a homogeneous medium, a result known in dissipative quantum mechanics [10].

The eigenstates of the BdG equation is written in the general form [8]

$$\psi_\alpha(x, y, z) = \frac{1}{\sqrt{L}} e^{ik_z z} \begin{pmatrix} e^{i(\mu - \frac{1}{2})\theta} u_\alpha(r) \\ e^{i(\mu + \frac{1}{2})\theta} v_\alpha(r) \end{pmatrix} \quad (6)$$

in the presence of a gap function $\Delta(\vec{r}) = \Delta(r)e^{-i\theta}$ for a singly-quantized vortex centered at the origin. The radial functions are obtained by solving the coupled differential equation

$$\begin{aligned} r^2 u'' + ru' + [(k_r^2 + E)r^2 - (\mu - 1/2)^2]u &= r^2 \Delta(r)v \\ r^2 v'' + rv' + [(k_r^2 - E)r^2 - (\mu + 1/2)^2]v &= -r^2 \Delta(r)u, \end{aligned} \quad (7)$$

with $k_r^2 = k_f^2 - k_z^2$. We set $2m \equiv 1$ for convenience and restore it in the final expression of M_v .

In arriving at the mass formula we made a physically important assumption that the dynamics of a vortex is *local in time*. From Eqs. (1) and (3) one finds that such assumption is valid if the vortex motion is *slow*, i.e. $r_v(\tau) \approx r_v(\tau')$, on the relevant time scale $|\tau - \tau'|$. For situations where this assumption is violated, one must work with the original, non-local action.

The eigenstates of the BdG equations are either *localized* at the core ($|E| < \Delta_\infty$), or *extended* ($|E| > \Delta_\infty$), where Δ_∞ is a gap at infinity. Accordingly the transition elements can be classified as *core-core*, *core-extended* (*extended-core*) and *extended-extended*. We examine the three contributions individually. The core solutions were first obtained by Caroli *et al.* [8] and the extended states are discussed by Cleary and by Bardeen *et al.* [11]

Core-core: The core states are given by

$$\begin{pmatrix} u(r) \\ v(r) \end{pmatrix} = \frac{1}{2} \left(\frac{k_r}{\xi_0} \right)^{\frac{1}{2}} \begin{pmatrix} J_{\mu - \frac{1}{2}}(k_r r) \\ J_{\mu + \frac{1}{2}}(k_r r) \end{pmatrix}. \quad (8)$$

The state is normalized over the radius ξ_0 . The coherence length ξ_0 is much larger than k_r^{-1} in extreme type-II superconductors considered here. Thus the derivatives of

the radial functions are essentially those of the Bessel functions, allowing us to apply the usual Bessel identities. We obtain

$$-\langle \alpha | \nabla_v H_0 | \beta \rangle = \frac{k_r \epsilon_0}{2} (\hat{x} \mp i\hat{y}) \delta_{\mu_\alpha, \mu_\beta \pm 1}, \quad (9)$$

where $\epsilon_0 = \Delta_\infty / k_r \xi_0$ is the core level spacing and $E_\alpha = \mu_\alpha \epsilon_0$. From this we obtain the mass:

$$\begin{aligned} M_v^{cc} &= m \left(\frac{4k_f L}{3\pi} \right) \left(\frac{E_f}{\epsilon_0} \right) \sum_\mu |f(\epsilon_0 \mu) - f(\epsilon_0 \mu + \epsilon_0)| \\ &\approx m \left(\frac{4k_f L}{3\pi} \right) \left(\frac{E_f}{\epsilon_0} \right) [f(-\Delta_\infty) - f(\Delta_\infty)]. \end{aligned} \quad (10)$$

The factor $k_f L$ counts the number of electrons along the length of the vortex L , given that the electron spacing is k_f^{-1} . Since $E_f / \epsilon_0 \sim (k_f \xi_0)^2$, the mass we obtained above is comparable to the mass of the electrons displaced by a hollow cylinder of radius ξ_0 . The zero temperature limit of Eq. (10) has been obtained previously [5,6]. The temperature dependence is through the difference of the Fermi distribution function at $\pm \Delta_\infty$, which is nearly constant for temperatures small compared with T_c . Close to T_c , the gap itself is sensitive to temperature, and Eq. (10) diverges as $\xi_0 \sim (T_c - T)^{-1/2}$.

This somewhat unphysical divergence of mass hints to us that perhaps our formula should not be taken too seriously near the critical temperature. One possible reason is the transition element, Eq. (9), which is based on Bessel function eigenstates. For diverging coherence length, it is doubtful that the expression (8) can be taken seriously all the way to $r = \xi_0 \gg k_r^{-1}$.

The spectral function for the core is a delta peak at frequency $\omega = \epsilon_0$. The (unapproximated) action due to the core-core transitions is

$$S_{eff} = \frac{k_f^3 \epsilon_0^2 L}{12\pi} \int_0^\beta d\tau \int_{-\infty}^\infty d\tau' |r_v(\tau) - r_v(\tau')|^2 e^{-\epsilon_0 |\tau - \tau'|}. \quad (11)$$

When $\epsilon_0 \sim (T_c - T)^{1/2}$ is small, the exponential factor $e^{-\epsilon_0 |\tau - \tau'|}$ remains order unity for a wide range of $|\tau - \tau'|$. For a large time interval the vortex may have moved a significant distance and not necessarily in a rectilinear fashion. In this case we cannot make the approximation leading to Eq. (3). It still remains true, however, that Eq. (11) governs the dynamics of a vortex. A motion described by Eq. (11) does not in general conserve energy. Hence a *fast* motion whose characteristic time scale is much smaller than ϵ_0^{-1} causes core quasiparticle transitions which dissipate energy.

Core-extended: We choose α for the extended states, and β for the core states. For the extended states, there are two independent solutions for a given energy, whose approximate forms are [11]

$$\begin{pmatrix} u(r) \\ v(r) \end{pmatrix}_{E > \Delta_\infty} = \sqrt{\frac{k_+}{2R}} \begin{pmatrix} u(E) \\ v(E) \end{pmatrix} \times J_{\mu-\frac{1}{2}}(k_+r), \\ \sqrt{\frac{k_-}{2R}} \begin{pmatrix} v(E) \\ u(E) \end{pmatrix} \times J_{\mu+\frac{1}{2}}(k_-r), \quad (12)$$

where $u(E), v(E) = \{(E \pm \sqrt{E^2 - \Delta^2(r)})/2E\}^{1/2}$ and $k_\pm = (k_r^2 \pm \sqrt{E^2 - \Delta^2(r)})^{1/2}$. The normalization is over a large cylinder of radius R . The negative energy solutions can be found from the substitution $(-v^*, u^*)$. Employing Eq. (5) for the mass,

$$|\langle \alpha | \nabla_v | \beta \rangle|^2 \approx \frac{\xi_0}{8R} \left(k_r - \frac{|\mu_\alpha|}{\xi_0} \right)^2 \theta(\xi_0 k_r - |\mu_\alpha|), \quad (13)$$

together with the constraint that the overlap integral is significant only if $\sqrt{E_\alpha^2 - \Delta_\infty^2} < \Delta_\infty$ [12]. The effective mass becomes ($x \equiv \beta \Delta_\infty / 2$)

$$M_v^{ce} = m \left(\frac{k_f L}{6\pi^2} \right) \left(\frac{E_f}{\epsilon_0} \right) \int_{-1}^1 d\mu \frac{(1-|\mu|)^\kappa}{1-\mu} \frac{\sinh[x(1-\mu)]}{\cosh[x] \cosh[x\mu]} \quad (14)$$

with $\kappa \approx 2.4$.

Replacing the integral with a characteristic value at $\mu = 0$ immediately shows that the core-extended vortex mass has the temperature dependence $\sim f(-\Delta_\infty) - f(\Delta_\infty)$, similar to the core-core mass. We conclude that M_v^{ce} is of the same order as M_v^{cc} , with the similar temperature dependence. The extended-core transitions will give the same mass as M_v^{ce} .

Extended-extended: The matrix element appearing in Eq. (4) consists of two parts. One involves the effect of “bending” of the order parameter $d\Delta(r)/dr$ and is only significant in the core region, $0 < r < \xi_0$. The other transition comes from the phase gradient $\nabla e^{-i\theta}$ which falls off as $1/r$, a long-ranged effect. One should also distinguish the transitions between states which lie on the same side of the Fermi surface, from those where a transition across the energy gap is involved. Their contributions to mass turn out to be very different.

We consider the $E_\alpha E_\beta > 0$ case first. Our approximate wavefunctions Eq. (12) and their negative energy counterparts are supposed to be valid for r not much bigger than ξ_0 . As r gets well outside the core the wavefunction picks up an additional phase shift not shown in Eq. (12) [11]. Directly substituting our wavefunctions in the phase-gradient part of the integral leads to very small values for $E_\alpha \approx E_\beta$, where the wavefunctions are likely to have the largest overlap. Hence a more accurate estimation of the wavefunction will not greatly enhance the size of the phase-gradient integral. We will thus assume that the integral can be restricted to the core region, where one can put $\Delta(r) = \Delta_\infty r / \xi_0$ [13]. With this substitution one has $-\langle \alpha | \nabla_v H | \beta \rangle \approx 2\pi \Delta_\infty \xi_0^{-1} (\hat{x} \mp i\hat{y}) \int_0^{\xi_0} r u_{\alpha(\beta)} v_{\beta(\alpha)} dr$. We need to separately consider four cases where α and β are one of the

extended eigenstates, and sum the squared magnitude of the transition amplitude for each process. We have

$$|\langle \alpha | \nabla_v H_0 | \beta \rangle|^2 \approx \frac{\Delta_\infty^2}{R^2} \theta(\xi_0 k_r - |\mu_\alpha|) \quad (15)$$

provided $(E_\alpha^2 - \Delta_\infty^2)^{1/2} + (E_\beta^2 - \Delta_\infty^2)^{1/2}$ is on the order of Δ_∞ or less. Both states lie close to the energy gap, reflecting the fact that states far from the Fermi surface do not influence the superconducting phenomenon.

We can compute the spectral function $J(\omega)$ according to Eq. (2). The sum over states is with respect to k_z , $\mu_\alpha = \mu_\beta \pm 1$, and the radial wavenumbers for both states. The integration measure is

$$\sum_{k_z} \sum_{\mu_\alpha \mu_\beta} \sum_{k_\alpha k_\beta} \approx \frac{LR^2 \xi_0^2}{2\pi^2} \iint \frac{|E_\alpha E_\beta| dE_\alpha dE_\beta}{\sqrt{(E_\alpha^2 - \Delta_\infty^2)(E_\beta^2 - \Delta_\infty^2)}}. \quad (16)$$

When $\omega = 0$, the integral is logarithmically divergent because of $(E^2 - \Delta_\infty^2)^{-1}$ term. Finite ω effectively serves as the lower cutoff, giving

$$J(\omega) \approx \frac{Lm^2 \xi_0 \Delta_\infty^2 (\Delta_\infty + \omega)}{\pi} \ln \left(\frac{\Delta_\infty}{\omega} \right) \times [f(\Delta_\infty) - f(\Delta_\infty + \omega)]. \quad (17)$$

In a previous calculation by two of the authors (Ao & Zhu), it was shown that a vortex experiences friction at non-zero temperature [7]. The friction coefficient η is obtained as the derivative of $J(\omega)$ at zero frequency [3]:

$$\eta = m \Delta_\infty \left(\frac{k_f L}{4\pi} \right) \frac{\beta \Delta_\infty / 2}{\cosh^2[\beta \Delta_\infty / 2]} \ln \left(\frac{\Delta_\infty}{\omega_c} \right) \quad (18)$$

with some lower frequency cutoff ω_c inserted by hand. A constant friction coefficient is a consequence of the *Ohmic* character of the spectral function, $J(\omega) \sim \omega$ (up to a logarithmic factor). In turn one finds from Eq. (3) that the mass in this case is linearly divergent! A straightforward evaluation of the mass gives

$$M_v^{ee} = m \left(\frac{k_f L}{2\pi^2} \right) \frac{\Delta_\infty}{\omega_c} \frac{\beta \Delta_\infty / 2}{\cosh^2[\beta \Delta_\infty / 2]} \ln \left(\frac{\Delta_\infty}{\omega_c} \right). \quad (19)$$

As anticipated, the mass diverges linearly for a fixed temperature as the low frequency cutoff tends to zero. For a fixed cutoff ω_c one gets exponentially vanishing mass in the zero temperature limit, because only the gapped transitions across the Fermi surface contribute to $J(\omega)$.

It was already mentioned that the very concept of mass depends on the motion the object executes. Here the mass is manifestly cutoff-dependent. To understand the physical origin of the cutoff ω_c let us go back to the original action, Eq. (1). One can divide the integral over ω into two parts; $0 < \omega < \omega_c$ and $\omega_c < \omega < \infty$, where a given

motion of a vortex is considered slow/fast in comparison with ω_c^{-1} . For the fast spectral mode in $\omega_c < \omega < \infty$ one can make the approximation $r_v(\tau) - r_v(\tau') \approx (\tau - \tau') \dot{r}_v(\tau)$ while for the more slowly responding environment one needs to keep the action as it is. If ω_c is much smaller than Δ_∞ , $J(\omega)$ is nearly Ohmic for $\omega < \omega_c$, and the action becomes

$$S_{eff} \approx \frac{1}{2} \int_0^\beta d\tau \left\{ \int_{\omega_c}^\infty d\omega \frac{4J(\omega)}{\pi\omega^3} \right\} \left(\frac{dr_v}{d\tau} \right)^2 + \frac{\eta}{2\pi} \int_0^\beta d\tau \int_{-\infty}^\infty d\tau' \left| \frac{r_v(\tau) - r_v(\tau')}{\tau - \tau'} \right|^2. \quad (20)$$

This is the standard Caldeira-Leggett model of a Brownian particle [3], where the mass and friction are given by Eqs. (19) and (18). The friction coefficient is related to the mass by $M_v^{ee} = 2\eta/\pi\omega_c$. The classical equation of motion becomes

$$M_v^{ee} \dot{v} + \eta v = \eta \left(\frac{2}{\pi\omega_c} \dot{v} + v \right). \quad (21)$$

Since the scale ω_c is itself set by the motion, it is reasonable to assume that $|\dot{v}| \approx |\omega_c v|$, indicating that the effects of inertia and drag are always of the same order regardless of the characteristic time scale of the motion.

Transitions across the Fermi level obviously do not contribute to friction, since the energy difference of the initial and final states is greater than $2\Delta_\infty$. We also expect that the mass contribution from this process is finite. In estimating the overlap integral we ignored the integration in the far region, $\xi_0 < r < R$, because the phase shifts for states E_α and E_β , $E_\alpha + E_\beta \approx 0$, should generically be different. Integrating over the core region gives

$$|\langle \alpha | \nabla_v H_0 | \beta \rangle|^2 \approx \frac{\Delta_\infty^2}{R^2} \left(2 - \frac{\Delta_\infty^2}{E_\alpha^2} \right) \quad (22)$$

provided $k_r \xi_0 > |\mu_\alpha| \approx |\mu_\beta|$. It is similar in magnitude to Eq. (15) for the $E_\alpha E_\beta > 0$ case. However the energy denominator is now of order $(E_\alpha - E_\beta)^3 \approx \Delta_\infty^3$ and free of divergence. The mass is given by $m(k_f L)[f(-\Delta_\infty) - f(\Delta_\infty)]$ up to a numerical factor of order unity. While the temperature dependence is similar to M_v^{cc} and M_v^{ce} numerically it is much smaller than both.

A vortex is not by itself assigned an intrinsic mass. Rather it is a consequence of interaction with the surrounding electrons. The motion of a vortex causes transitions between various BdG eigenstates and accrues mass as a consequence.

We showed that both M_v^{cc} and $M_v^{ce(ec)}$ are comparable to the mass of electrons displaced by the hollow core in the $T = 0$ limit, and has a temperature dependence $\sim \xi_0^2(T)[f(-\Delta_\infty) - f(\Delta_\infty)]$. The transition between extended states with small energy differences leads to a

frequency-dependent mass and a friction at finite temperature. Both quantities vanish at zero temperature exponentially. Considering extended states alone, one concludes that the vortex motion is dissipative at finite T even in the absence of impurities. In the low temperature range where M_v^{cc} and M_v^{ce} are large and the friction is small, the vortex motion is Newtonian (mass times acceleration equals force), with a large mass. Experimental indications may have already existed [14]. If the characteristic time scale of the motion is much smaller than the inverse core level spacing, however, the core level transitions can lead to dissipation also.

P. Ao and X.-M. Zhu were supported by the Swedish Natural Science Research Council (NFR).

-
- [1] J. Bardeen and M. J. Stephen, Phys. Rev. **140**, A1197 (1965); P. Nozières and W. F. Vinen, Phil. Mag. **14**, 667 (1966).
 - [2] G. Blatter, V. B. Geshkenbein, and V. M. Vinokur, Phys. Rev. Lett. **66**, 3297 (1991); P. Ao and D. J. Thouless, Phys. Rev. Lett. **72**, 132 (1994).
 - [3] A. O. Caldeira and A. J. Leggett, Ann. Phys. (NY) **149**, 374 (1983).
 - [4] H. Suhl, Phys. Rev. Lett. **14**, 226 (1965); J.-M. Duan and A. J. Leggett, Phys. Rev. Lett. **68**, 1216 (1992).
 - [5] N. B. Kopnin and M. M. Salomaa, Phys. Rev. B **44**, 9667 (1991); T. C. Hsu, Physica C **213**, 305 (1993); D. M. Gaitonde and T. V. Ramakrishnan, Phys. Rev. B **56**, 11951 (1997); N. B. Kopnin and V. M. Vinokur, cond-mat/9806065.
 - [6] E. Šimánek, Phys. Lett. A **194**, 323 (1994).
 - [7] P. Ao and X.-M. Zhu, Physica C **282-287**, 367 (1997).
 - [8] P. G. de Gennes, *Superconductivity of Metals and Alloys* (Addison-Wesley, 1989); C. Caroli, P. G. de Gennes, and J. Matricon, Phys. Lett. **9**, 307 (1964).
 - [9] We consider a rectilinear vortex of length L along the z direction.
 - [10] V. Hakim and V. Ambegaokar, Phys. Rev. A **32**, 423 (1985).
 - [11] R. M. Cleary, Phys. Rev. **175**, 587 (1968); J. Bardeen, R. Kümmel, A. E. Jacobs, and L. Tewordt, Phys. Rev. **187**, 556 (1969).
 - [12] The constraint follows from integrating over a product of two Bessel functions whose arguments are $k_r r$ and $k_\pm r$, respectively, over the coherence length, since $|k_r - k_\pm|$ differing by more than ξ_0^{-1} will make the integral very small.
 - [13] F. Gygi and M. Schluter, Phys. Rev. B **43**, 7609 (1991); N. Hayashi *et al.*, Phys. Rev. Lett. **80**, 2921 (1998).
 - [14] G.T. Seidler *et al.*, Phys. Rev. Lett. **74**, 1442 (1995).